

Prospects for observing CP violation and rare decays at ATLAS and CMS

Maria Smizanska
Lancaster University, UK

For ATLAS and CMS collaborations

Investigating B-hadron decays represents an alternative approach to direct searches for physics beyond the Standard Model (BSM). ATLAS and CMS concentrate on B decays that can be registered by a di-muon signature. B-hadrons decaying to $J/\psi \rightarrow \mu^+\mu^-$ will statistically dominate B-physics analyses allowing high precision measurements, in particular a test of BSM effects in the CP violation of $B_s^0 \rightarrow J/\psi\phi$. In the so-called rare B-decay sector, ATLAS and CMS will concentrate on a family of semi-muonic exclusive channels, $b \rightarrow s\mu^+\mu^-$, and on the purely muonic decay $B_s^0 \rightarrow \mu^+\mu^-$. After three years of LHC running at a luminosity of a few times $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ (corresponding to 30 fb^{-1}), each of these two experiments can measure the $B_s^0 \rightarrow \mu^+\mu^-$ signal with 3σ significance, assuming the Standard Model (SM) value for the decay probability.

I. INTRODUCTION

In the past decade BaBar, Belle, CDF and D0 made very precise measurements of flavour and CP-violating phenomena. Whilst the analysis of the remaining data of these experiments may still push the boundaries, no evidence of physics beyond the SM, nor any evidence for CP violation other than that originating from the CKM mechanism, has yet been found. At LHC, thanks to the large beauty production cross-section and the high luminosity of the machine, the sensitivity of B-decay measurements is expected to substantially improve. Whilst direct detection of new particles in ATLAS and CMS will be the main approach to discovering new physics, these two experiments will also perform B-Physics searches to provide complementary information on BSM physics from B-decays. In the present document we concentrate on three of the most prominent cases: the rare decay $B_s^0 \rightarrow \mu^+\mu^-$, CP violation in $B_s^0 \rightarrow J/\psi\phi$ and the semi-muonic exclusive channels of type $b \rightarrow s\mu^+\mu^-$.

II. TRIGGERS FOR B-DECAYS WITH DI-MUON SIGNATURES

Both the ATLAS and CMS triggers comprise three levels and the selection of B-decays with two muons in the final state is initiated by a di-muon at the first trigger level (L1). At luminosity $10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ the lowest possible threshold of about $p_T > 4 \text{ GeV}$ (3 GeV) in ATLAS (CMS) will be used, rising to 6 - 8 GeV at $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. The muon is confirmed at the second trigger level (L2) combining muon and inner detector (ID) information. In addition this trigger makes cuts on the invariant mass and certain secondary vertex qualities of the B decay products to select specific physics processes. Figure 1 illustrates the $B_s^0 \rightarrow \mu^+\mu^-$ mass resolution and the $J/\psi \rightarrow \mu^+\mu^-$ vertex reconstruction in the CMS and ATLAS L2 triggers respectively. The channels $B_s^0 \rightarrow J/\psi\phi$ and $B_s^0 \rightarrow \mu^+\mu^-$ are triggered by requiring di-muons fulfilling J/ψ or B_s^0 mass cuts. For semi-muonic rare channels, such as $B_d \rightarrow K^{*\circ}\mu^+\mu^-$ or $B_s \rightarrow \phi\mu^+\mu^-$, ID tracks are combined to first reconstruct the $K^{*\circ}$ or ϕ ; the muon tracks are then added to reconstruct the B. Selected events are further analysed at the third trigger level by algorithms using full event information. Approximately ten B-physics events will be recorded per second to permanent storage in each ATLAS and CMS.

III. SENSITIVITY IN $B_s^0 \rightarrow \mu^+\mu^-$

In the SM the di-muonic rare decay $B_s^0 \rightarrow \mu^+\mu^-$ is mediated by flavour-changing neutral currents that are forbidden at tree level. In addition this decay is helicity suppressed, resulting in an expected branching ratio (BR)




FIG. 1: Reconstruction of di-muon signals in the second-level trigger. CMS $B_s^0 \rightarrow \mu^+\mu^-$ mass reconstruction using fast L2 track reconstruction (left), and the same for the offline reconstruction (middle). Right, L2 di-muon vertex fit (χ^2) for the $J/\psi \rightarrow \mu^+\mu^-$ signal and for di-muon background in ATLAS.




FIG. 2: Upper limits on the branching ratio of $B_s^0 \rightarrow \mu^+\mu^-$ for each ATLAS and CMS. The results are given at confidence level of 90% as a function of integrated luminosity.

of $(3.42 \pm 0.52) \times 10^{-9}$ [1]. The motivation for precise measurements of rare B-decays is to test the SM up to high perturbative orders, as well as to search for BSM effects that can increase the BR compared to SM predictions. After $B_s^0 \rightarrow \mu^+\mu^-$ events have passed the triggers the offline selection criteria are applied, see Table I. With 30 fb^{-1} (expected after three years of running at a luminosity of a times of $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$), ATLAS and CMS will be able to select 17.1 ± 6 and 18.3 ± 6.3 signal events respectively, assuming the SM value for the BR. Number of 42^{+39}_{-30} (ATLAS) and 42^{+66}_{-42} (CMS) background events $b\bar{b} \rightarrow \mu^+\mu^- X$ are expected. The errors are dominated by limited statistics of simulated events. Contributions from other backgrounds: $B_s^0 \rightarrow K^-\mu^+\nu$, $B_d^0 \rightarrow K^-\pi^+$ and $B_d^0 \rightarrow \pi^-\pi^+$ are negligible, see [2] and [3] for reference. These results were used to estimate upper limits on the branching ratio of $B_s^0 \rightarrow \mu^+\mu^-$, shown in Figure 2. Both ATLAS and CMS were using the algorithms of Ref. [4]. After three years at luminosity $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$, both ATLAS and CMS can reconstruct the $B_s^0 \rightarrow \mu^+\mu^-$ signal with significance 3σ . The strategy is to continue the di-muon channel programme up to nominal LHC luminosity $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$.

IV. CP VIOLATION AND OTHER PARAMETERS OF $B_s^0 \rightarrow J/\psi\phi$ DECAY

New phenomena beyond the SM may alter CP violation in B-decays. A prominent example that has received experimental and theoretical attention is the decay $B_s^0 \rightarrow J/\psi\phi$. Combined D0 and CDF results on a pair of parameters – the mixing phase ϕ_s and the decay width difference $\Delta\Gamma_s$ – yields a 2.2σ deviation from the SM [5]. The LHC is expected to produce the statistics necessary to unambiguously confirm or reject possible BSM

ATLAS			CMS		
Selection cut	$B_s^0 \rightarrow \mu^+ \mu^-$	$b\bar{b} \rightarrow \mu^+ \mu^- X$	Selection cut	$B_s^0 \rightarrow \mu^+ \mu^-$	$b\bar{b} \rightarrow \mu^+ \mu^- X$
$I_{\mu\mu}^{atl} > 0.9$, $L_{xy} > 0.5\text{mm}$, $\alpha < 0.017$ rad, $-70 \text{ MeV} < \Delta m < 140 \text{ MeV}$	$\epsilon = 0.04$	$\epsilon = 0.24 \cdot 10^{-6}$	$I_{\mu\mu}^{cms} > 0.85$, $L_{xy} > 0.8\text{mm}$ $\alpha < 0.1$ rad $ \Delta m < 100 \text{ MeV}$	$\epsilon = 0.016$	$\epsilon = 0.26 \cdot 10^{-6}$
Event yield 30 fb^{-1}	17.1 ± 6	42_{-30}^{+39}		18.3 ± 6.3	42_{-42}^{+66}

TABLE I: Selection efficiencies, ϵ , and numbers of signal and background events for 30 fb^{-1} . Efficiencies were calculated after preselection criteria had been applied: $p_T^\mu > 6(4)\text{GeV}$ for the first and second muon (ATLAS), $p_T^\mu > 3\text{GeV}$ and $p_T^{\mu\mu} > 5\text{GeV}$ (CMS); $4 \text{ GeV} < m(\mu\mu) < 7.3 \text{ GeV}$ (ATLAS) and $5 \text{ GeV} < m(\mu\mu) < 6 \text{ GeV}$ (CMS). Δm denotes a mass difference $m(\mu\mu) - M_{B_s}$. Variables $I_{\mu\mu}^{cms}$, $I_{\mu\mu}^{atl}$ of di-muon isolation, as well as the angle α and the distance L_{xy} have been defined in [2] and [3] for ATLAS and CMS respectively.

	ATLAS	CMS	Statistical error on parameter	ATLAS	CMS
Signal statistics, 30 fb^{-1}	$2.4 \cdot 10^5$	$2.6 \cdot 10^5$	$\delta\phi_s$	0.067	0.067
Proper time resolution	0.083 ps	0.077 ps	$\delta\Delta\Gamma_s/\Delta\Gamma_s$	13%	12%
B_s^0 mass resolution	16.6 MeV	14 MeV	$\delta\Gamma_s/\Gamma_s$	1%	0.9%
Background	$\sim 30\%$	$\sim 33\%$	$\delta A_{ }/A_{ }$	0.7%	0.8%
Tag quality using μ , e, jet-charge tags	3.9	3.9	$\delta A_{\perp}/A_{\perp}$	3%	2.7%

TABLE II: $B_s^0 \rightarrow J/\psi\phi$ – summary of performance and expected statistical errors of parameters for an integrated luminosity of 30 fb^{-1} . Background is dominated by $B_d^0 \rightarrow J/\psi K^{0*}$, $B_d^0 \rightarrow J/\psi K^+ \pi^-$ decays.

contributions in this decay. With an integrated luminosity of 30 fb^{-1} ATLAS and CMS will accumulate $2.4 \cdot 10^5$ and $2.6 \cdot 10^5$ $B_s^0 \rightarrow J/\psi\phi$ events respectively. Simulation of the time dependent three-dimensional angular distribution of the final state particles proved that both experiments can perform a model independent determination of ϕ_s and $\Delta\Gamma_s$, along with the average width Γ_s and some parameters of three helicity amplitudes of the decay. The factors determining the measurement sensitivity – the B_s^0 proper decay-time resolution, flavour tag quality, signal statistics and background – are summarized in Table II. Under these assumptions the weak phase ϕ_s will be determined with a precision of 0.067 in both ATLAS and CMS. Whilst this precision is not enough to confirm the SM estimate of ϕ_s (-0.0368 ± 0.0018), it could verify a possible enhancement from BSM contributions as predicted for instance in [6]. The expected precisions of other parameters are given in Table II. The strong phases of the helicity amplitudes have been found to be highly correlated in the fits and it is thus supposed that these will be determined from the process $B_d^0 \rightarrow J/\psi K^{0*}$.

V. PROGRAMME FOR RARE EXCLUSIVE SEMILEPTONIC DECAYS

In the SM, the electroweak penguin decays $b \rightarrow s(d)l^+l^-$ are only induced at the one-loop level, leading to small branching fractions and thus a sensitivity to contributions from BSM. Problematic hadronic form-factor uncertainties entering the calculations of branching ratios lead physicists to observables that are less unaffected by the uncertainties and are therefore a better basis for BSM searches. The most frequently used forward – backward asymmetry A_{FB} is defined through the angle between the μ^+ and the B-hadron momenta in di-muon rest frame [7]. In ATLAS the shape of the A_{FB} distribution as a function of the di-muon mass is negligibly affected by applying di-muon triggers (Fig. 3). This feature will make the results only weakly dependent on trigger acceptance calibrations [2]. A list of decays supported by ATLAS is given in Table III together with the corresponding statistics after offline selections have reduced the background. The background determination was limited by MC statistics and therefore only an upper limit is given. Sensitivity studies were performed for each channel and in particular for Λ_b^0 decay a precision of 6% on the A_{FB} can be achieved after 30 fb^{-1} , see Fig. 3. This will allow to distinguish the SM from one example

Process	Signal	$b\bar{b} \rightarrow \mu^+\mu^- X$	Process	Signal	$b\bar{b} \rightarrow \mu^+\mu^- X$
$B_d^0 \rightarrow \mu^+\mu^- K^{0*}$	2500	< 10000	$B_s^0 \rightarrow \mu^+\mu^- \phi$	900	< 10000
$B^+ \rightarrow \mu^+\mu^- K^+$	2300	< 10000	$\Lambda_b^0 \rightarrow \mu^+\mu^- \Lambda^0$	800	< 4000
$B^+ \rightarrow \mu^+\mu^- K^{*+}(K_S^0\pi^+)$	4000	< 10000			

TABLE III: Signal and background statistics in semileptonic rare decays in ATLAS.

of MSSM, calculated for $\Lambda_b^0 \rightarrow \mu^+\mu^- \Lambda^0$ in Ref. [8]



FIG. 3: Left, A_{FB} for $B_s^0 \rightarrow \mu^+\mu^- \phi$ events in ATLAS: before L1 trigger (filled triangles), after di-muon L1 trigger with $p_T^\mu > 4$ GeV (filled circles) and with $p_T^\mu > 6$ GeV (open squares). Right, ATLAS sensitivity of A_{FB} measurement in $\Lambda_b^0 \rightarrow \mu^+\mu^- \Lambda^0$, three large points with error bars show ATLAS sensitivity with 30 fb^{-1} , upper set of points generated by SM, lower set by MSSM model in Ref. [8].

VI. CONCLUSIONS

The ATLAS and CMS trigger and offline tools for di-muon B-physics channels are well prepared for the luminosities up to few times of $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. Both experiments are ready to perform measurements which will be sufficiently precise to allow confirmation of some possible BSM physics contributions.

-
- [1] G. Buchalla and A. J. Buras, Nucl. Phys. B400, 225 (1993); A.J. Buras, Phys. Lett. B566, 115 (2003).
 - [2] ATLAS Collaboration, "Expected Performance of the ATLAS Experiment, Detector, Trigger and Physics", CERN-OPEN-2008-020, Geneva, 2008, to appear.
 - [3] C. Eggel, U. Langenegger and A. Starodumov, CMS-CR-2006-071.
 - [4] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
 - [5] D. Strom for CDF and D0 collaborations, "Bs parameters in $B_s^0 \rightarrow J/\psi \phi$ ", current proceedings.
 - [6] P. Ball "Probing new physics through B(s) mixing", 42nd Rencontres de Moriond on Electroweak Interactions and Unified Theories, La Thuile, 10-17 Mar 2007, e-Print: hep-ph/0703214.
 - [7] C. Bobeth, G. Hiller, G. Piranishvili, JHEP 0712:040 (2007).
 - [8] T. M. Aliev, A. Ozpineci and M. Savci, Nucl.Phys.B 649 (2003) 168.